

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 November 2002 (21.11.2002)

PCT

(10) International Publication Number  
WO 02/092761 A2

- (51) International Patent Classification<sup>7</sup>: C12N
- (21) International Application Number: PCT/US02/14570
- (22) International Filing Date: 7 May 2002 (07.05.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/851,673 8 May 2001 (08.05.2001) US
- (71) Applicant (for all designated States except US): IMMUNEX CORPORATION [US/US]; Law Department, 51 University Street, Seattle, WA 98101 (US).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

- (72) Inventors; and
- (75) Inventors/Applicants (for US only): DERRY, Jonathan, M.J. [GB/US]; 3998 El Cimo Lane, Bainbridge Island, WA 98110 (US). FANSLOW, William, C., III [US/US]; 404 S.W. 197th Street, Normandy Park, WA 98166 (US). DOUGALL, William, Conrad [US/US]; 7523 33rd Avenue Northwest, Seattle, WA 98117 (US).
- (74) Agent: PERKINS, Patricia, Anne; IMMUNEX CORPORATION, 51 University Street, Seattle, WA 98101 (US).

(54) Title: SCREENING ASSAYS FOR AGONISTS OR ANTAGONISTS OF CD40 SIGNALING



(57) Abstract: The present invention provides methods for identifying a molecule that antagonizes or agonizes CD40 activity by screening for molecules that modulate the binding of NEMO and CYLD. Also provided are assays useful in screening for the binding of NEMO and CYLD and databases and methods useful for analyzing the resultant information.



WO 02/092761 A2

## Screening Assays for Agonists or Antagonists of CD40 Signaling

### Field of the Invention

This invention relates to methods for screening for agonists and/or antagonists of activities associated with TNF receptor family member CD40.

### Background of the Invention

CD40 is a member of the tumor necrosis factor (TNF)/nerve growth factor (NGF) receptor family, which is defined by the presence of cysteine-rich motifs in the extracellular region (Smith et al., *Science* 248:1019, 1990; Mallett and Barclay, *Immunology Today* 12:220, 1991; Locksley et al., *Cell* 104:487, 2001). This family includes the lymphocyte antigen CD27, CD30 (an antigen found on Hodgkin's lymphoma and Reed-Sternberg cells), two receptors for TNF, a murine protein referred to as 4-1BB, rat OX40 antigen, NGF receptor, and *Fas* antigen. Similar to other members of the TNF/NGF receptor family, CD40 contains a leader sequence, trans-membrane domain an extracellular domain responsible for binding its native cognate, CD40 ligand (CD40L; described in U.S. Patent Nos. 5,961,974, 5,962,406 and 5,981,724; hereinafter, the Armitage patents). CD40 has been found to be expressed on B lymphocytes, monocytes/macrophages, dendritic cells, smooth muscle cells, microglia, epithelial cells and some carcinoma cell lines.

The cytoplasmic domain of CD40 associates intracellularly with several of the TNF receptor-associated factors (TRAFs; Baker and Reddy, *Oncogene* 12:1 (1996)) including TRAFs 2, 3, 5 and 6 (Pullen et al., *Biochemistry* 37:11836, 1998; Cao et al., *Nature* 383:443, 1996; Ishida et al., *Proc. Natl. Acad. Sci. USA* 93:9437, 1996). The TRAFs are cytoplasmic proteins that often mediate signal transduction by members of the TNF receptor superfamily, and they are important in the regulation of, for example, immune and inflammatory responses.

Triggering of CD40, such as by contact with membrane-bound or soluble CD40 ligand (CD40L), results in the stimulation of CD40-mediated cellular responses. These cellular responses can include the activation of transcription factor NF-kappaB, a ubiquitous transcription factor that is extensively utilized in cells of the immune system. NF-kappaB is normally maintained in the cytoplasm by interaction with IkappaB members (reviewed in Karin, *Oncogene* 18: 6867, 1999; Chen and Gosh, *Oncogene*

18:6845, 1999). Release of NF-kappaB, and subsequent translocation of this transcription factor to the nucleus, result from phosphorylation of IkappaB, which leads to its ubiquitination and degradation. Phosphorylation of IkappaB is mediated by the inducible kappaB kinase (IKK) complex consisting of two kinases, IKKalpha, and  
5 IKKbeta, and a scaffolding protein referred to as NEMO (NF-kappaB essential modulator; Yamaoka et al., *Cell* 93:1231, 1998), IKK-gamma (Rothwarf et al., *Nature* 395:297, 1998), FIP3 (Fourteen K interacting protein 3; Li et al., *Proc. Natl. Acad. Sci. USA* 96:1042, 1999) or RAP2 (RIP-associated protein 2; WO 99/47672) (hereinafter, NEMO).

10 NEMO exhibits characteristics of a "signaling scaffold protein", and binds not only to kinases involved in NF-kappaB-mediated signaling, but to proteins that play a role in signaling via TNF and/or Fas (Receptor Interacting Protein 2 or RIP2, McCarty et al., *J. Biol. Chem.* 273:16968, 1998; NF-kappaB Inducing Kinase, NIK, Malinin et al., *Nature* 385:640, 1997; and A20, Cooper et al., *J. Biol. Chem.* 271:18068, 1996), to several viral  
15 proteins (Adeno 14.7K, Li et al., *supra*; HTLV Tax, Yamaoka et al., *supra*), and to proteins of unknown function (TIP60, Tat interactive protein 60 kDa, Kamine et al., *Virology* 216:357, 1996; CYLD, a putative tumor suppressor gene associated with familial cylindromatosis, Bignell et al., *Nat. Genet.* 25:160, 2000).

Additionally, a peptide that blocks the interaction of NEMO with IKK-alpha/IKK-  
20 beta has been shown to be potent anti-inflammatory agent (May et al., *Science* 289:1550, 2000). Moreover, deletion mutations of NEMO have been shown to cause familial incontinentia pigmentosa (IP; Smahi et al., *Nature* 405:466, 2000), whereas milder mutations in the carboxy-terminal region of NEMO have been linked to ectodermal dysplasia with immunodeficiency/hyperIgM (HED-ID; Zonano et al., *Am. J. Hum. Genet.*  
25 67:1555, 2000; Jain et al., *Nature Immunology* 2:223, 2001). Because of the role NEMO plays in numerous signaling pathways, there is a need in the art to identify molecules that can affect the interaction of NEMO with certain molecules (either as antagonists or agonists), to allow regulation of specific pathways in an inflammatory response.

#### Summary of the Invention

30 The present invention provides methods for screening for a molecule that antagonizes or agonizes the activity of NEMO in CD40 signaling. In one aspect of the invention, there is provided a method for identifying compounds that alter CD40

signaling activity comprising: (a) mixing a test compound with a polypeptide selected from the group consisting of (i) a NEMO polypeptide comprising amino acids 287 through 419 SEQ ID NO:2, (ii) a fragment of a NEMO polypeptide according to (i) that is capable of binding a CYLD polypeptide according to SEQ ID NO:4 or fragment or variant thereof, and (iii) variants of the NEMO polypeptides of (i) and (ii); and (b) determining whether the test compound alters the ability of NEMO to bind CYLD. The invention further provides a method for identifying compounds that inhibit binding of NEMO and CYLD comprising: (a) mixing a test compound with a polypeptide selected from the group consisting of (i) a NEMO polypeptide comprising amino acids 287 through 419 SEQ ID NO:2, (ii) a fragment of a NEMO polypeptide according to (i) that is capable of binding a CYLD polypeptide according to SEQ ID NO:4, and (iii) variants of the NEMO polypeptides of (i) and (ii), and a binding partner of said NEMO polypeptide selected from the group consisting of (iv) a CYLD polypeptide according to SEQ ID NO:4, (v) a fragment of a CYLD polypeptide according to SEQ ID NO:4 that is capable of binding a NEMO polypeptide of (i), (ii), or (iii), and (vi) variants of the CYLD polypeptides of (iv) and (v); and (b) determining whether the test compound inhibits the binding activity of said NEMO and CYLD polypeptides.

In one aspect, the inventive methods utilize homogeneous assay formats such as fluorescence resonance energy transfer, fluorescence polarization, time-resolved fluorescence resonance energy transfer, scintillation proximity assays, reporter gene assays, fluorescence quenched enzyme substrate, chromogenic enzyme substrate and electrochemiluminescence. In another aspect, the inventive methods utilize heterogeneous assay formats such as enzyme-linked immunosorbant assays (ELISA) or radioimmunoassays. In yet another aspect of the invention are cell-based assays, for example those utilizing reporter genes, as well as functional assays that analyze the effect of an antagonist or agonist on biological function(s).

The invention further provides methods for producing information comprising the identity of a compound that alters one or more biological activities of CD40, the method comprising using assay methods of the invention to identify one or more compounds that alter the binding of NEMO and CYLD. In one preferred embodiment, the compound decreases (or antagonizes) the binding of NEMO and CYLD, and in another distinct embodiment, the compound increases (or agonizes) the binding of NEMO and CYLD.

Preferably the biological activity of CD40 that is decreased or downregulated is selected from the group consisting of deleterious effects of CD40-mediated immune or inflammatory response (including atherosclerosis, arthritis, multiple sclerosis (MS), systemic lupus erythematosus (SLE), thrombosis, graft versus host disease and/or graft rejection. In a distinct embodiment, the biological activity of CD40 that is increased or upregulated is selected from the group consisting of upregulation of a cell-mediated immune response, upregulation of an antibody-mediated immune response, prevention or treatment of infectious disease, and prevention or treatment of neoplastic disease.

Also provided by the invention is the information produced according to the inventive methods, said information comprising the identity of a compound that alters the biological activity of CD40, and preferably embodied in a storage medium selected from the group consisting of the brains of living organisms, paper, magnetic tape, optical tape, floppy disks, compact disks, computer system hard drives, and computer memory units. In a further aspect, the invention provides a database comprising said information, wherein the information is preferably embodied in a computer-readable medium, and a separate embodiment wherein the information is embodied in a human-readable medium.

Additionally provided by the invention is a computer system comprising a database containing records pertaining to a plurality of compounds, wherein the records comprise results of an assay of the invention, and a user interface allowing a user to access information regarding the plurality of compounds. In another aspect of the invention, a computer system is provided for storing and retrieving data on a plurality of compounds, the computer system comprising: (a) input means for entering data for the compounds into a storage medium; (b) a processor for creating an individual record for each compound, the processor assigning specific identifying values for each compound; (c) means for selecting one or more of the records based on results in an assay; and (d) means for transmitting information in the record or records to an output device to produce a report; preferably a report in human-readable form, and wherein the computer system preferably further comprises a video display unit.

The invention also provides a method of using the computer system of the invention to select one or more compounds for testing from a plurality of compounds having records stored in a database, the method comprising: displaying a list of said records or a field for entering information identifying one or more of said records; and selecting one or more of the records from the list or the record or records identified by

entering information in the field. Further, the invention provides a method of operating a computer system for analyzing compounds that modulate the interaction NEMO and CYLD, the method comprising: (a) entering data relating to a plurality of compounds into a storage medium; (b) processing the data to create an individual record for each compound; (c) testing compounds for the ability to modulate binding of NEMO to CYLD; and (d) communicating results from the testing into the storage medium such that results for each compound are associated with the individual record for that compound; wherein in one embodiment the storage medium comprises one or more computer memory units, and in another embodiment the computer system further comprises a video display unit.

In yet another aspect of the invention, a database is provided comprising records generated according to the methods of the invention, and a method is provided for selecting compounds that modulate the interaction of NEMO and CYLD, comprising compiling said database, analyzing the testing results, and selecting one or more compounds.

Candidate molecules that are determined to agonize or antagonize a CD40 signaling activity of NEMO are useful, for example, for the further definition of CD40-mediated signaling pathways, and for the manipulation of CD40-mediated cellular responses. Moreover, CD40 signaling agonists and antagonists provide therapeutic agents for treating disorders of the immune system, and inflammatory disorders, as well as treatment of conditions characterized by malignant cells expressing CD40.

#### Brief Description of the Drawings

Figure 1 depicts the structure NEMO, with various regions highlighted; the amino acid sequence is shown in SEQ ID NO:2. The region from amino acid 217 through 419 is believed to bind CYLD; homodimerization is thought to be mediated by amino acids 217 through 264. Binding to NIK and TIP60 is believed to occur between amino acids 95 and 264, while binding to RIP and A20 is through amino acids 95 through 218. NEMO binds to 14.7K via amino acids 180 through 419; binding to IKK-1 and IKK-2 is mediated by amino acids 44 through 86. The majority of mutations that result in familial IP are deletions of exons 3 through 10. The stippled regions labeled ' $\alpha$ H' are alpha-helical regions 'LZ' represents the leucine zipper region, and 'ZF' denotes the zinc finger. Amino acids 397, 400, 413 and 417 are believed to coordinate zinc.

### Detailed Description of the Preferred Embodiment

The present invention provides methods for screening for a molecule that antagonizes or agonizes CD40 signaling activity, utilizing molecules that play a role in signaling through CD40. Two patients with a form of X-linked hyper IgM syndrome with ectodermal dysplasia and normal CD40 and CD40L expression were found to have mutations in NEMO. In one, a T to C mutation at nucleotide 1249 results in a Cys to Arg substitution at amino acid 417 of NEMO (C417R), and in the other, an A to T substitution at nucleotide 1217 results in an Asp to Val mutation at amino acid 406 (V406D). B cells from these patients did not undergo immunoglobulin class switching when contacted with a soluble, oligomeric form of CD40L (described in the Armitage patents, *supra*), and also failed to upregulate CD54 expression. Moreover, antigen-presenting cells from these patient did not synthesize Interleukin-12 (IL-12) and TNF-alpha when stimulated with CD40L and interferon-gamma (IFN-gamma), but did synthesize these two cytokines when stimulated with *S. aureus* Cowan protein A plus IFN-gamma or lipopolysaccharide plus IFN-gamma. Additionally, degradation of IkappaB-alpha in response to TNF or LPS appeared normal in monocytes from these patients, but stimulation with CD40L failed to induce this response.

These findings demonstrated that the zinc finger region of NEMO represents an area of NEMO that is critical for CD40 signal transduction, but that does not appear critical for transduction of signal via other pathways leading to NF-kappaB activation (for example, binding of LPS through the Toll-like receptors, and/or binding of TNF to TNF receptor). Accordingly, peptides derived from this region are likely to be useful in screening for small molecules that inhibit the interaction of this domain of NEMO with other molecules in the CD40 signaling cascade (i.e., CYLD), which will be useful in downregulating or controlling deleterious effects of CD40-mediated immune or inflammatory response. Conditions that are thought to be mediated by CD40 signaling include atherosclerosis, arthritis, multiple sclerosis (MS), systemic lupus erythematosous (SLE), thrombosis, graft versus host disease and/or graft rejection.

#### Candidate Molecules to be Tested for CD40 Signaling Activity:

The methods of the invention may be used to identify antagonists and agonists of CD40 signaling activity from cells, cell-free preparations, chemical libraries, cDNA libraries, recombinant antibody libraries (or libraries comprising subunits of antibodies)

and natural product mixtures. The antagonists and agonists may be natural or modified substrates, ligands, enzymes, receptors, etc. of the polypeptides of the instant invention, or may be structural or functional mimetics of one of the polypeptides (NEMO or CYLD). Potential antagonists of the instant invention may include small molecules, peptides and antibodies that bind to and occupy a binding site of the inventive polypeptides or a binding partner thereof, causing them to be unavailable to bind to their natural binding partners and therefore preventing normal biological activity. Potential agonists include small molecules, peptides and antibodies which bind to the instant polypeptides or binding partners thereof, and elicit the same or enhanced biologic effects as those caused by the binding of the polypeptides of the instant invention.

Small molecule agonists and antagonists are usually less than 10K molecular weight and may possess a number of physicochemical and pharmacological properties which enhance cell penetration, resist degradation and prolong their physiological half-lives (Gibbs, J., *Pharmaceutical Research in Molecular Oncology*, Cell, Vol. 79 (1994)). Antibodies, which include intact molecules as well as fragments such as Fab and F(ab')<sub>2</sub> fragments, as well as recombinant molecules derived therefrom (including antibodies expressed on phage, intrabodies, single chain antibodies such as scFv and other molecules derived from immunoglobulins that are known in the art), may be used to bind to and inhibit the polypeptides of the instant invention by blocking the propagation of a signaling cascade. It is preferable that the antibodies are humanized, and more preferable that the antibodies are human. The antibodies of the present invention may be prepared by any of a variety of well-known methods.

Additional examples of candidate molecules, also referred to herein as "test molecules," to be tested for CD40 signaling agonist or antagonist activity include, but are not limited to, carbohydrates, small molecules (usually organic molecules or peptides), proteins, and nucleic acid molecules (including oligonucleotide fragments typically consisting of from 8 to 30 nucleic acid residues). Peptides to be tested typically consist of from 5 to 25 amino acid residues. Also, candidate nucleic acid molecules can be antisense nucleic acid sequences, and/or can possess ribozyme activity.

Small molecules to be screened using the hereindescribed screening assays can typically be administered orally or by injection to a patient in need thereof. Small molecules that can be administered orally are especially preferred. The small molecules of the invention preferably will not be toxic at the doses required for them to be effective



as pharmaceutical agents, and they are preferably not subject to rapid loss of activity in the body, such as the loss of activity that might result from rapid enzymatic or chemical degradation. In addition, pharmaceutically useful small molecules are preferably not immunogenic.

5           The methods of the invention can be used to screen for antisense molecules that inhibit the functional expression of one or more mRNA molecules that encode one or more proteins that mediate a CD40-dependent cellular response. An anti-sense nucleic acid molecule is a DNA sequence that is inverted relative to its normal orientation for transcription and so expresses an RNA transcript that is complementary to a target mRNA  
10 molecule expressed within the host cell (*i.e.*, the RNA transcript of the anti-sense nucleic acid molecule can hybridize to the target mRNA molecule through Watson-Crick base pairing). An anti-sense nucleic acid molecule may be constructed in a number of different ways provided that it is capable of interfering with the expression of a target protein. Typical anti-sense oligonucleotides to be screened preferably are 30-40  
15 nucleotides in length. The anti-sense nucleic acid molecule generally will be substantially identical (although in antisense orientation) to the target gene. The minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity  
20 would be most preferred.

Candidate nucleic acid molecules can possess ribozyme activity. Thus, the methods of the invention can be used to screen for ribozyme molecules that inhibit the functional expression of one or more mRNA molecules that encode one or more proteins that mediate a CD40 dependent cellular response. Ribozymes are catalytic RNA  
25 molecules that can cleave nucleic acid molecules having a sequence that is completely or partially homologous to the sequence of the ribozyme. It is possible to design ribozyme transgenes that encode RNA ribozymes that specifically pair with a target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself  
30 altered, and is thus capable of recycling and cleaving other molecules. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the antisense constructs.

The design and use of target RNA-specific ribozymes is described in Haseloff et al. (*Nature*, 334:585, 1988; see also U.S. Patent No.5,646,023), both of which publications are incorporated herein by reference. Tabler et al. (*Gene* 108:175, 1991) have greatly simplified the construction of catalytic RNAs by combining the advantages of the anti-sense RNA and the ribozyme technologies in a single construct. Smaller regions of homology are required for ribozyme catalysis, therefore this can promote the repression of different members of a large gene family if the cleavage sites are conserved.

#### NEMO and CYLD Molecules

10 Generally, the screening assays described herein involve a NEMO and a CYLD protein, or nucleic acid encoding such. NEMO and the NEMO-binding molecule CYLD and nucleic acids encoding these proteins are known in the art. The nucleotide sequence of a DNA encoding NEMO, and amino acid sequence encoded by this DNA, are set forth in SEQ ID NOs:1 and 2, respectively. The nucleotide sequence of a DNA encoding  
15 CYLD, and amino acid sequence encoded by this DNA, are set forth in SEQ ID NOs:3 and 4, respectively. However, it is understood that other NEMO and CYLD variants other than those shown in these examples may be used in the hereindisclosed assays, including other NEMO and CYLD molecules known in the art, or variants having similar properties.

20 Sequence variants of native NEMO and CYLD polypeptides are useful in the practice of the present invention in any instance where the native NEMO or CYLD polypeptide is utilized, provided that the variant possesses any biological activity required for the assay. Generally for these assays, suitable NEMO variants will bind CYLD. Mutations present in such variants may include, for example, substitutions, deletions, and  
25 insertions of amino acids. Allelic forms or mutated forms of NEMO and CYLD can be obtained for use in these assays by using a variety of techniques known in the art, including, for example, site-directed mutagenesis, oligonucleotide-directed mutagenesis, and so on.

Also useful in the inventive methods are fragments of NEMO and/or CYLD.  
30 Particularly useful fragments of NEMO include the region from about amino acid 300 to 419, comprising a leucine zipper and zinc finger domain, and the region from about amino acid 387 to 419, comprising the zinc finger domain (see Figure 1); additional fragments thereof that bind CYLD can be identified as described herein, and will also be

useful in the present methods. Such fragments include those that are truncated by about five to ten amino acids (i.e., fragments from x to y, wherein x is selected from the group consisting of 386, 385, 384, 3843, 382, 381, 380 379, 378 and 377, and y is selected from the group consisting of 409, 410, 411, 412, 413, 414, 415, 416, 417, 418 and 419, and in particular, 418 and 419), and those having an N-terminus between amino acid 300 and 387 (i.e., fragments from x to y, wherein x is an integer between 300 and 387, and y is selected from the group consisting of 409, 410, 411, 412, 413, 414, 415, 416, 417, 418 and 419). Particularly useful fragments of CYLD include those that are capable of binding NEMO.

Sequence variants of NEMO and CYLD polypeptides that are not capable of binding their native binding partner may also be useful in any of the assays described herein. In one embodiment, such sequence variants are useful as controls for an assay. In another embodiment, sequence variants that do not bind their respective binding partners will be useful in screening for molecules that facilitate the binding of NEMO and CYLD despite the inability of the sequence variants to bind in the absence of the facilitating molecule. Such facilitating molecules will be useful in treating disease conditions characterized by an inability of NEMO to bind to CYLD (i.e., X-linked hyper IgM with ectodermal dysplasia). Useful sequence variants can be obtained as described herein.

NEMO and/or CYLD peptides that are useful in the inventive methods (including fragments such as those mentioned previously) may be expressed as fusion proteins with tag peptides that facilitate detection and/or purification. Such peptides include, for example, poly-His or the antigenic identification peptides described in U.S. Patent No. 5,011,912 and in Hopp et al., *Bio/Technology* 6:1204, 1988. Additional, useful tag proteins include green fluorescent protein (GFP; Chalfie et al., *Science* 263:802, 1994), an N-terminal peptide that contains recognition sites for a monoclonal antibody, a specific endopeptidase, and a site-specific protein kinase (PKA; Blonar and Rutter, *Science* 256:1014, 1992), birA (Altman et al., *Science* 274:94, 1996), and glutathione S transferase (GST; Smith and Johnson, *Gene* 67:31, 1988).

One such tag peptide is the FLAG<sup>TM</sup> peptide, which is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. A murine hybridoma designated 4E11 produces a monoclonal antibody that binds the FLAG<sup>TM</sup> peptide in the presence of certain divalent metal cations, as described in U.S. Patent 5,011,912, hereby

incorporated by reference. The 4E11 hybridoma cell line has been deposited with the American Type Culture Collection under accession no. HB 9259. Monoclonal antibodies that bind the FLAG<sup>TM</sup> peptide are available from Eastman Kodak Co., Scientific Imaging Systems Division, New Haven, Connecticut.

5 Another useful tag peptide is the GST peptide, which binds glutathione, also facilitating purification of expressed recombinant protein. Recombinant protein can be purified by affinity chromatography using a suitable chromatography matrix to which has been attached glutathione, as described in Smith and Johnson, supra, hereby incorporated by reference. Suitable chromatography matrixes include Glutathione-Agarose beads  
10 (Pharmacia, Uppsala, Sweden). Recombinant protein can be eluted with an excess of glutathione. Fragments of NEMO comprising the zinc finger domain (as described above) and a GST peptide are preferred.

The proteins useful in the practice of the present invention typically have an amino acid sequence that is at least 80% identical, or at least 85% identical, or preferably  
15 at least 90% identical to all or a portion of the corresponding native protein as set forth in SEQ ID NOS:2 or 4. Percent identity is defined as the percentage of the amino acid residues set forth in SEQ ID NOS:2, or 4, that are identical with part or all of another protein sequence (which may be a portion of a larger protein sequence) after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent identity.  
20 For comparing amino acid sequences of unequal length, the percent identity is calculated based on the smaller of the two sequences.

Preferably, the comparison is done using a computer program. An exemplary, preferred computer program is the Genetics Computer Group (GCG; Madison, WI) Wisconsin package version 10.0 program, 'GAP.' The preferred default parameters for  
25 the 'GAP' program includes: (1) The GCG implementation of the previously stated comparison matrixes for nucleotides and amino acids; (2) a penalty of 30 for each gap and an additional penalty of 1 for each symbol in each gap for amino acid sequences, or penalty of 50 for each gap and an additional penalty of 3 for each symbol in each gap for nucleotide sequences; (3) no penalty for end gaps; and (4) no maximum penalty for long  
30 gaps. Another program useful for determining percent identify is the BESTFIT program, also available from the University of Wisconsin as part of the GCG computer package. Default parameters for using the BESTFIT program are the same as those described above for using the GAP program.

Screening Assays for NEMO and CYLD:

Specific screening methods are known in the art and along with integrated robotic systems and collections of chemical compounds/natural products are extensively incorporated in high throughput screening so that large numbers of test compounds can be tested for antagonist or agonist activity within a short amount of time. These methods include homogeneous assay formats such as fluorescence resonance energy transfer, fluorescence polarization, time-resolved fluorescence resonance energy transfer, scintillation proximity assays, reporter gene assays, fluorescence quenched enzyme substrate, chromogenic enzyme substrate and electrochemiluminescence, as well as more traditional heterogeneous assay formats such as enzyme-linked immunosorbant assays (ELISA) or radioimmunoassays. Homogeneous assays are preferred. Also comprehended herein are cell-based assays, for example those utilizing reporter genes, as well as functional assays that analyze the effect of an antagonist or agonist on biological function(s) (for example, secretion of cytokines or immunoglobulin class switching).

Moreover, combinations of screening assays can be used to find molecules that regulate the biological activity of NEMO and/or CYLD. Molecules that regulate the biological activity of a polypeptide may be useful as agonists or antagonists of the peptide. In using combinations of various assays, it is usually first determined whether a candidate molecule binds to a polypeptide by using an assay that is amenable to high throughput screening. Binding candidate molecules identified in this manner are then added to a biological assay to determine biological effects. Molecules that bind and that have an agonistic or antagonistic effect on biologic activity will be useful in treating or preventing disease or conditions with which the polypeptide(s) are implicated.

Generally, an antagonist will inhibit the activity by at least 30%; more preferably, antagonists will inhibit activity by at least 50%, most preferably by at least 90%. Similarly, an agonist will enhance the activity by at least 20%; more preferably, agonists will enhance activity by at least 30%, most preferably by at least 50%. Those of skill in the art will recognize that agonists and/or antagonists with different levels of agonism or antagonism respectively may be useful for different applications (i.e., for treatment of different disease states).

Homogeneous assays are mix-and-read style assays that are very amenable to robotic application, whereas heterogeneous assays require separation of free from bound

analyte by more complex unit operations such as filtration, centrifugation or washing. These assays are utilized to detect a wide variety of specific biomolecular interactions (including protein-protein, receptor-ligand, enzyme-substrate, and so on), and the inhibition thereof by small organic molecules. These assay methods and techniques are well known in the art (see, e.g., High Throughput Screening: The Discovery of Bioactive Substances, John P. Devlin (ed.), Marcel Dekker, New York, 1997 ISBN: 0-8247-0067-8). The screening assays of the present invention are amenable to high throughput screening of chemical libraries and are suitable for the identification of small molecule drug candidates, antibodies, peptides, and other antagonists and/or agonists, natural or synthetic.

One such assay is based on fluorescence resonance energy transfer (FRET; for example, HTRF®, Packard BioScience Company, Meriden, CT; LANCE™, PerkinElmer LifeSciences, Wallac Oy., Turku, Finland) between two fluorescent labels, an energy donating long-lived chelate label and a short-lived organic acceptor. The energy transfer occurs when the two labels are brought in close proximity via the molecular interaction between NEMO and CYLD. In a FRET assay for detecting inhibition of the binding of NEMO and CYLD, europium chelate or cryptate labeled NEMO or CYLD serves as an energy donor and streptavidin-labeled allophycocyanin (APC) bound to the appropriate binding partner (i.e., CYLD if NEMO is labeled, or NEMO if CYLD is labeled) serves as an energy acceptor. Once NEMO binds CYLD, the donor and acceptor molecules are brought in close proximity, and energy transfer occurs, generating a fluorescent signal at 665 nm. Antagonists of the interaction of NEMO and CYLD will thus inhibit the fluorescent signal, whereas agonists of this interaction would enhance it.

Another useful assay is a bioluminescence resonance energy transfer, or BRET, assay, substantially as described in Xu et al., *Proc. Natl. Acad. Sci. USA* 96:151 (1999). Similar to a FRET assay, BRET is based on energy transfer from a bioluminescent donor to a fluorescent acceptor protein. However, a green fluorescent protein (GFP) is used as the acceptor molecule, eliminating the need for an excitation light source. Exemplary BRET assays include BRET and BRET<sup>2</sup> from Packard BioScience, Meriden, CT.

DELFLIA® (dissociated enhanced lanthanide fluoroimmunoassay; PerkinElmer LifeSciences, Wallac Oy., Turku, Finland) is a solid-phase assay based on time-resolved fluorometry analysis of lanthanide chelates (see, for example, US Patent 4,565,790, issued January 21, 1986). For this type of assay, microwell plates are coated with a first

protein (NEMO or CYLD). The binding partner (CYLD or NEMO, respectively) is conjugated to europium chelate or cryptate, and added to the plates. After suitable incubation, the plates are washed and a solution that dissociates europium ions from solid phase bound protein, into solution, to form highly fluorescent chelates with ligands present in the solution, after which the plates are read using a reader such as a VICTOR<sup>2</sup>™ (PerkinElmer LifeSciences, Wallac Oy., Turku, Finland) plate reader to detect emission at 615 nm).

Another assay that will be useful in the inventive methods is a FlashPlate® (Packard Instrument Company, IL)-based assay. This assay measures the ability of compounds to inhibit protein-protein interactions. FlashPlates® are coated with a first protein (either NEMO or CYLD), then washed to remove excess protein. For the assay, compounds to be tested are incubated with the second protein (CYLD, if the plates are coated with NEMO, or NEMO if plates are coated with CYLD) and I<sup>125</sup> labeled antibody against the second protein and added to the plates. After suitable incubation and washing, the amount of radioactivity bound is measured using a scintillation counter (such as a MicroBeta® counter; PerkinElmer LifeSciences, Wallac Oy., Turku, Finland).

The AlphaScreen™ assay (Packard Instrument Company, Meriden, CT). AlphaScreen™ technology is an "Amplified Luminescent Proximity Homogeneous Assay" method utilizing latex microbeads (250 nm diameter) containing a photosensitizer (donor beads), or chemiluminescent groups and fluorescent acceptor molecules (acceptor beads). Upon illumination with laser light at 680 nm, the photosensitizer in the donor bead converts ambient oxygen to singlet-state oxygen. The excited singlet-state oxygen molecules diffuse approximately 250 nm (one bead diameter) before rapidly decaying. If the acceptor bead is in close proximity to the donor bead (i.e., by virtue of the interaction of NEMO and CYLD), the singlet-state oxygen molecules reacts with chemiluminescent groups in the acceptor beads, which immediately transfer energy to fluorescent acceptors in the same bead. These fluorescent acceptors shift the emission wavelength to 520-620 nm, resulting in a detectable signal. Antagonists of the interaction of NEMO and CYLD will thus inhibit the shift in emission wavelength, whereas agonists of this interaction would enhance it.

One embodiment of a method for identifying molecules which inhibit or antagonize CD40-mediated signaling involves adding a candidate molecule to a medium which contains cells that express NEMO and CYLD; changing the conditions of said

medium so that, but for the presence of the candidate molecule, NEMO would be bound to CYLD, and observing the binding and stimulation or inhibition of a functional response. The activity of the cells that were contacted with the candidate molecule may then be compared with the identical cells that were not contacted and antagonists and agonists of the polypeptides of the instant invention may be identified. The measurement of biological activity may be performed by a number of well-known methods such as measuring the amount of protein present (e.g. an ELISA) or of the protein's activity. A decrease in biological stimulation or activation would indicate an antagonist. An increase would indicate an agonist.

5

#### Computer Analysis of Assay Results

10

15

20

In one aspect of the invention, the assays of the invention are used to identify compounds that alter CD40 signaling activity. The benefits of integrated robotic systems used to analyze collections of chemical compounds/natural products in such assays, which preferably incorporate high-throughput screening methods, are most often realized by the use of sophisticated computer and statistical techniques to manage the resulting data. In one form, the information generated in the inventive screening assays is stored (or compiled) in electronic form, using a computerized database that allows information to be efficiently catalogued and retrieved. Such databases are comprised of records, usually one record for each compound, that includes information about the compound, such as chemical name, structure, source, activity in a binding assay, activity in a biological assay, etc.

25

30

The information may be entered into the database manually, that is by a user entering data through a user interface (i.e., keyboard, touchpad, etc.), or it may be entered electronically as in when a robotic system for analysis of compounds generates electronic results that are transferred to another computer system (often referred to as uploading). Such information is usually stored in a discrete area of the record referred to as a field. Additionally, the information, preferably in the form of a database, may be stored permanently or temporarily on various forms of storage media, including paper, the brains of living organisms, compact disks, floppy disks, magnetic tapes, optical tapes, hard drives, computer system memory units, and the like.

The database may be stand-alone, or the records therein may be related to other databases (a relational database). Examples of other databases include publicly available,



well-known databases such as GenBank for peptides and nucleic acids (and associated databases maintained by the National Center for Biotechnology Information or NCBI), and the databases available through [www.chemfinder.com](http://www.chemfinder.com) or The Dialog Corporation (Cary, North Carolina) for chemical compounds.

5 A user will be able to search the database according to the information recorded (selecting records that have a particular value in a selected field, for example, searching for all compounds that inhibited a binding assay by at least about 30%); accordingly, another aspect of the invention is a method of using a computer system to catalog and store information about various chemical compounds. The ability to store and retrieve  
10 such information in computerized form allows those of ordinary skill in the art to select compounds for additional testing, including additional analysis of binding ability, biological testing, and testing in animal models or clinical trials of pharmaceutical agents in humans. Moreover, in addition to storing and cataloging information, the database can be used to provide a report, either in electronic form or in the form of a printout, that will  
15 facilitate further analysis of selected compounds.

One embodiment of the invention comprises a computing environment; an input device, connected to the computing environment, to receive information from the user; an output device, connected to the computing environment, to provide information to the user; and a plurality of algorithms selectively executed based on at least a portion of the  
20 received information, wherein any one of these algorithms analyzes at least a portion of the received information and generates output information, and preferably wherein the output information is communicated via the output device. The computing environment preferably further comprises a communications network; a server connected to the network; and a client connected to the network, wherein the client is part of a client-  
25 server architecture and typically is an application that runs on a personal computer or workstation and relies on a server to perform some operations (see Nath, 1995, *The Guide To SQL Server*, 2nd ed., Addison-Wesley Publishing Co.).

The computing environment of the present invention is advantageously implemented using any multipurpose computer system including those generally referred  
30 to as personal computers and mini-computers. Such a computer system will include means for processing input information such as at least one central processor, for example an Intel® processor (including Pentium® Pentium® II, Celeron™, Pentium® II3,

Pentium® 4 or the like), or Motorola processor (for example, a PowerPC G3 or PowerPC G4 microprocessor capable of running at speeds up to 533 MHz or higher); a storage device, such as a hard disk, for storing information related to CD40, NEMO and/or CYLD polypeptides and/or compounds that alter the binding of NEMO and CYLD (or signaling through CD40); and means for receiving input information. Those of skill in the art recognize that computer technology is changing at a rapid rate; accordingly, new, improved versions of processors are comprehended herein.

The processor, which comprises and/or accesses memory units of the computer system, is programmed to perform analyses of information related to the CD40, NEMO and/or CYLD polypeptides and/or compounds that modulate the binding of NEMO and CYLD (or signaling through CD40). This programming may be permanent, as in the case where the processor is a dedicated PROM (programmable read-only memory) or EEPROM (electrically erasable programmable read-only memory), or it may be transient in which case the programming instructions are loaded from the storage device or from a floppy diskette or other transportable computer-readable media. The computing environment further preferably comprises a user interface such as a Unix/X-Window interface, a Microsoft Windows interface, or a Macintosh operating system interface.

Preferably, the computing environment further includes an optical disk for storing data, a printer for providing a hard copy of the data, and a monitor or video display unit to facilitate user input of information and to display both input and output information. The output information may be output from the processor within the computer system in print form using a printer; on a video display unit; or via a communications link or network to another processor or client application.

The following examples are intended to illustrate various embodiments of the invention, and should not be construed to limit the invention.

#### EXAMPLE 1

This example describes a gene promoter/reporter system based on the human Interleukin-8 (IL-8) promoter used to analyze the activation of gene transcription in vivo. Other NF-kappaB-responsive promoters besides IL-8 could be used including a minimal promoter element comprising NF-kappaB consensus binding sites. The induction of human IL-8 gene transcription by the cytokines Interleukin-1 (IL-1) or tumor necrosis

factor-alpha (TNF- $\alpha$ ) is known to be dependent upon intact NF-kappaB and NF-IL-6 transcription factor binding sites. Fusion of the cytokine-responsive IL-8 promoter with a cDNA encoding the murine IL-4 receptor (mIL-4R) allows measurement of promoter activation by detection of the heterologous reporter protein (mIL-4R) on the cell surface of transfected cells. Other detectable moieties may be used, including any protein or peptide that is detectable by a selected assay and is not present on or in the cells under other conditions (for example, luciferase or human IL-2 receptor).

Human kidney epithelial cells (293/EBNA) are transfected (via the DEAE/DEXTRAN method) with a plasmid encoding the reporter/promoter construct (referred to as pIL-8rep), and cultured under conditions promoting viability. If CD40 activation is necessary for NEMO/CYLD association, 293 cells are transfected with plasmids encoding CD40. Alternatively, the NF-kappaB responsive promoter is introduced (by transfection) into cells that express endogenous CD40 (for example, 70Z/3, WEHI-231, or RAW 264.7 cell-lines). Stimulation of CD40 may be accomplished via addition of soluble CD40L, agonistic antibodies to CD40, or by cotransfection with the transmembrane form of CD40L; use of soluble CD40L is preferred.

The transfected cells are contacted with compounds to be tested. If a compound binds NEMO or CYLD, and inhibits the interaction of these two proteins, NF-kappaB will remain sequestered and unable to activate the IL-8 promoter. As a result, there will be no mIL-4R present on the surface of cells that have been contacted with an effective antagonist. The presence or absence of the mIL-4 receptor is detected by a radioimmunoassay (RIA) or other suitable assay.

## EXAMPLE 2

This example illustrates the association of NEMO with CYLD. Interaction of NEMO with CYLD is demonstrated by co-immunoprecipitation assays essentially as described by Hsu et al. (*Cell* 84:299; 1996). Briefly, 293/EBNA cells are co-transfected with plasmids that direct the synthesis of NEMO and epitope-tagged CYLD (or CYLD and epitope-tagged NEMO). Other cells may also be used, including cells that constitutively express NEMO and/or CYLD (for example, 70Z/3 cells). Two days after transfection, cells are lysed (for example, in a buffer containing 0.5% NP-40 and labeled (for example, with biotin-ester). NEMO and proteins associated with NEMO are immunoprecipitated with anti-NEMO (or anti-CYLD), washed extensively, resolved by

electrophoretic separation on a 6-10% SDS polyacrylamide gel and electrophoretically transferred to a nitrocellulose membrane for Western blotting. The association of NEMO with CYLD is visualized by probing the membrane with an antibody that specifically recognizes the tag.

5

### EXAMPLE 3

This example describes a yeast two-hybrid screening assay that is useful in screening compounds for the ability to modulate the binding of NEMO and CYLD. Yeast comprising (1) an expression cassette encoding a GAL4 DNA binding domain (or GAL4 activator domain) fused to a binding fragment of NEMO capable of binding to a  
10 GAL4 activator domain) fused to a binding fragment of NEMO capable of binding to a CYLD polypeptide, (2) an expression cassette encoding a GAL4 DNA activator domain (or GAL4 binding domain, respectively) fused to a binding fragment of CYLD capable of binding to a NEMO polypeptide, and (3) a reporter gene (e.g., beta-galactosidase) comprising a cis-linked GAL4 transcriptional response element can be used for agent  
15 screening. Such yeast are incubated with a test agent or the appropriate control(s) under conditions promoting expression of the reporter gene in the absence of an inhibitor, and expression of the reporter is determined. The capacity of the agent to modulate expression of the reporter gene as compared to a control culture identifies the agent as a candidate modulatory agent.

20

### EXAMPLE 4

This example illustrates the preparation of monoclonal antibodies against NEMO or CYLD. Preparations of purified NEMO or CYLD, for example, or transfected cells expressing high levels of NEMO or CYLD, are employed to generate monoclonal  
25 antibodies against NEMO or CYLD using conventional techniques, such as those disclosed in U.S. Pat. No. 4,411,993, incorporated herein by reference. DNA encoding NEMO or CYLD can also be used as an immunogen, for example, as reviewed by Pardoll and Beckerleg in Immunity 3: 165, 1995. Such antibodies are likely to be useful in interfering with NEMO/CYLD binding (antagonistic or blocking antibodies), as  
30 components of diagnostic or research assays for NEMO or CYLD activity, or in affinity purification of NEMO or CYLD.

To immunize rodents, NEMO or CYLD immunogen is emulsified in an adjuvant (such as complete or incomplete Freund's adjuvant, alum, or another adjuvant, such as

Ribi adjuvant R700 (Ribi, Hamilton, Mont.), and injected in amounts ranging from 10-100 µg subcutaneously into a selected rodent, for example, BALB/c mice or Lewis rats. DNA may be given intradermally (Raz et al., *Proc. Natl. Acad. Sci. USA* 91: 9519, 1994) or intramuscularly (Wang et al., *Proc. Natl. Acad. Sci. USA* 90: 4156, 1993); saline has  
5 been found to be a suitable diluent for DNA-based antigens. Ten days to three weeks days later, the immunized animals are boosted with additional immunogen and periodically boosted thereafter on a weekly, biweekly or every third week immunization schedule.

Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot-blot assay (antibody sandwich), ELISA (enzyme-linked immunosorbent  
10 assay), immunoprecipitation, or other suitable assays, including FACS analysis. Following detection of an appropriate antibody titer, positive animals are given an intravenous injection of antigen in saline. Three to four days later, the animals are sacrificed, splenocytes harvested, and fused to a murine myeloma cell line (e.g., NS1 or preferably Ag 8.653 [ATCC CRL 1580]). Hybridoma cell lines generated by this  
15 procedure are plated in multiple microtiter plates in a selective medium (for example, one containing hypoxanthine, aminopterin, and thymidine, or HAT) to inhibit proliferation of non-fused cells, myeloma-myeloma hybrids, and splenocyte-splenocyte hybrids.

Hybridoma clones thus generated can be screened by ELISA for reactivity with NEMO or CYLD, for example, by adaptations of the techniques disclosed by Engvall et  
20 al., *Immunochem.* 8: 871 (1971) and in U.S. Pat. No. 4,703,004. A preferred screening technique is the antibody capture technique described by Beckman et al., *J. Immunol.* 144: 4212 (1990). Positive clones are then injected into the peritoneal cavities of syngeneic rodents to produce ascites containing high concentrations (>1 mg/ml) of monoclonal antibody. The resulting monoclonal antibody can be purified by ammonium  
25 sulfate precipitation followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can also be used, as can affinity chromatography based upon binding to NEMO or CYLD protein.

Techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, *Science* 242:423, 1988; Huston et al., *Proc. Natl. Acad. Sci. USA*  
30 85:5879, 1988; and Ward et al., *Nature* 334:544, 1989) can also be adapted to produce single chain antibodies against NEMO or CYLD. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge,

resulting in a single chain polypeptide. Such single chain antibodies can be useful intracellularly (i.e., as 'intrabodies), for example as described by Marasco *et al.* (*J. Immunol. Methods* 231:223-238, 1999) for genetic therapy in HIV infection. Intrabodies that bind NEMO or CYLD and inhibit the interaction thereof will be useful in  
5 downregulating specific CD40-signaling response.

### EXAMPLE 5

In accordance with the present invention, a series of oligonucleotides are designed to target different regions of the NEMO or CYLD mRNA molecule, using the nucleotide  
10 sequence of SEQ ID Nos:1 or 3, respectively, as the basis for the design of the oligonucleotides. The oligonucleotides are selected to be approximately 10, 12, 15, 18, or more preferably 20 nucleotide residues in length, and to have a predicted hybridization temperature that is at least 37 degrees C. Preferably, the oligonucleotides are selected so that some will hybridize toward the 5' region of the mRNA molecule, others will  
15 hybridize to the coding region, and still others will hybridize to the 3' region of the mRNA molecule.

The oligonucleotides may be oligodeoxynucleotides, with phosphorothioate backbones (internucleoside linkages) throughout, or may have a variety of different types of internucleoside linkages. Generally, methods for the preparation, purification, and use  
20 of a variety of chemically modified oligonucleotides are described in U.S. Patent No. 5,948,680. As specific examples, the following types of nucleoside phosphoramidites may be used in oligonucleotide synthesis: deoxy and 2'-alkoxy amidites; 2'-fluoro amidites such as 2'-fluorodeoxyadenosine amidites, 2'-fluorodeoxyguanosine, 2'-fluorouridine, and 2'-fluorodeoxycytidine; 2'-O-(2-methoxyethyl)-modified amidites such as  
25 2,2'-anhydro[1-(beta-D-arabino-furanosyl)-5-methyluridine], 2'-O-methoxyethyl-5-methyluridine, 2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine, 3'-O-acetyl-2'-O-methoxy-ethyl-5'-O-dimethoxy-trityl-5-methyluridine, 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine, 2'-O-methoxyethyl-5'-O-dimethoxy-trityl-5-methylcytidine, N4-benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-  
30 cytidine, and N4-benzoyl-2'-O-methoxyethyl-5'-O-di-methoxytrityl-5-methylcytidine-3'-amidite; 2'-O-(aminoxyethyl) nucleoside amidites and 2'-O-(dimethylaminoxyethyl) nucleoside amidites such as 2'-(dimethylaminoxyethoxy) nucleoside amidites, 5'-O-tert-butyl-diphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine, 5'-O-tert-butyl-diphenylsilyl-2'-O-(2-

hydroxyethyl)-5-methyluridine, 2'-O-[(2-phthalimidoxy)ethyl]-5'-*t*-butyl-diphenyl-silyl-5-methyluridine, 5'-O-*tert*-butyl-diphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine, 5'-O-*tert*-butyl-diphenylsilyl-2'-O-[N,N-dimethyl-aminooxyethyl]-5-methyluridine, 2'-O-(dimethyl-aminoxy-ethyl)-5-methyluridine, 5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine, and 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropyl-phosphor-amidite]; and 2'-(aminooxyethoxy) nucleoside amidites such as N2-isobutyryl-6-O-diphenyl-carbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

10 Modified oligonucleosides may also be used in oligonucleotide synthesis, for example methylenemethylimino-linked oligonucleosides, also called MMI-linked oligonucleosides; methylene-dimethylhydrazo-linked oligonucleosides, also called MDH-linked oligonucleosides; methylene-carbonylamino-linked oligonucleosides, also called amide-3-linked oligonucleosides; and methylene-aminocarbonyl-linked oligonucleosides, also called amide-4-linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages, which are prepared as described in U.S. Pat. Nos. 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289. Formacetal- and thioformacetal-linked oligonucleosides may also be used and are prepared as described in U.S. Pat. Nos. 5,264,562 and 5,264,564; and ethylene oxide  
15 linked oligonucleosides may also be used and are prepared as described in U.S. Pat. No. 5,223,618. Peptide nucleic acids (PNAs) may be used as in the same manner as the oligonucleotides described above, and are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, Bioorganic & Medicinal Chemistry, 1996, 4, 5-23; and U.S. Pat.  
20 Nos. 5,539,082, 5,700,922, and 5,719,262.

Chimeric oligonucleotides, oligonucleosides, or mixed oligonucleotides-oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound.  
30 Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers". Some examples of different types of chimeric

oligonucleotides are: [2'-O-Me]--[2'-deoxy]--[2'-O-Me] chimeric phosphorothioate oligonucleotides, [2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides, and [2'-O-(2-methoxy-ethyl)phosphodiester]--[2'-deoxyphosphoro-thioate]--[2'-O-(2-methoxyethyl)-phosphodiester] chimeric oligonucleo-  
5 tides, all of which may be prepared according to U.S. Patent No. 5,948,680. In one preferred embodiment, chimeric oligonucleotides ("gapmers") 18 nucleotides in length are utilized, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside  
10 (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines. Other chimeric oligonucleotides, chimeric oligonucleosides, and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to U.S. Pat. No. 5,623,065.

Oligonucleotides are preferably synthesized via solid phase P(III)  
15 phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. The concentration of oligonucleotide in each well is assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products is evaluated by capillary electrophoresis, and base and backbone composition is confirmed by mass  
20 analysis of the compounds utilizing electrospray-mass spectroscopy.

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. Cells are routinely maintained for up to 10 passages as  
25 recommended by the supplier. When cells reach 80% to 90% confluency, they are treated with oligonucleotide. For cells grown in 96-well plates, wells are washed once with 200 microliters OPTI-MEM-1 reduced-serum medium (Gibco BRL) and then treated with 130 microliters of OPTI-MEM-1 containing 3.75 g/mL LIPOFECTIN (Gibco BRL) and the desired oligonucleotide at a final concentration of 150 nM. After 4 hours of treatment,  
30 the medium is replaced with fresh medium. Cells are harvested 16 hours after oligonucleotide treatment. Preferably, the effect of several different oligonucleotides should be tested simultaneously, where the oligonucleotides hybridize to different



portions of the target nucleic acid molecules, in order to identify the oligonucleotides producing the greatest degree of inhibition of expression of the target nucleic acid.

Antisense modulation of NEMO or CYLD nucleic acid expression can be assayed in a variety of ways known in the art. For example, NEMO or CYLD mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation and Northern blot analysis are taught in, for example, Ausubel, F. M. et al., Current Protocols in Molecular Biology, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1996.

Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, Calif. and used according to manufacturer's instructions. This fluorescence detection system allows high-throughput quantitation of PCR products. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE or FAM, obtained from either Operon Technologies Inc., Alameda, Calif. or PE-Applied Biosystems, Foster City, Calif.) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, Calif. or PE-Applied Biosystems, Foster City, Calif.) is attached to the 3' end of the probe.

When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular (six-second) intervals by laser optics built into the ABI PRISM 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from

untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples. Other methods of quantitative PCR analysis are also known in the art.

NEMO or CYLD protein levels can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA, or fluorescence-activated cell sorting (FACS). Antibodies directed to NEMO or CYLD polypeptides can be prepared via conventional antibody generation methods such as those described herein. Immunoprecipitation methods, Western blot (immunoblot) analysis, and enzyme-linked immunosorbent assays (ELISA) are standard in the art (see, for example, Ausubel, F. M. et al., Current Protocols in Molecular Biology, Volume 2, pp. 10.16.1-10.16.11, 10.8.1-10.8.21, and 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991).

#### EXAMPLE 6

This example describes a mammalian two-hybrid screening assay (substantially as described in Shioda et al., *Proc. Natl. Acad. Sci. USA* 97:5220, 2000) that is useful in screening compounds for the ability to modulate the binding of NEMO and CYLD. For this assay, CV-1/EBNA-1 monkey kidney epithelial cells expressing Epstein-Barr virus nuclear antigen 1 (EBNA-1) are stably transfected with a reporter plasmid for GAL4-dependent expression of the green fluorescent protein (GFP). Clones that express GFP when transfected transiently with transcriptional activators fused to GAL4 DNA-binding domain with minimal background GFP expression are identified.

Useful clones are then transfected stably with a model bait (for example, NEMO or CYLD) and prey (CYLD or NEMO, respectively); under conditions in which both NEMO and CYLD are expressed and allowed to bind to each other, the cells will express GFP and can be readily identified by green fluorescence in cell culture. Such cells are incubated with a test agent or the appropriate control(s) and the capacity of the agent to modulate expression of GFP as compared to a control culture is determined. Those of skill in the art will be able to select other reporter genes as desired.

#### EXAMPLE 7

In this assay, test molecules capable of modulating an activity associated with CD40 signaling are identified by assessing the ability of B cells to undergo immunoglobulin class switching, and/or the ability of peripheral blood mononuclear cells

(PBMCs) to synthesize IL-12 and/or TNF-alpha. Human peripheral blood mononuclear cells (PBMC) are isolated from peripheral blood from normal volunteers by density gradient centrifugation over Histopaque<sup>®</sup> (Sigma, St. Louis, MO). T cell-depleted preparations of cells (E<sup>-</sup>) are obtained by removing T cells by rosetting with 2-aminoethylisothiuronium bromide-treated SRBC (sheep red blood cells) and further  
5 density gradient centrifugation over Histopaque<sup>®</sup>. Alternatively, B cells are selectively isolated using anti CD19-coated magnetic beads (Dynal, Lake Success, NY).

B cell assays are conducted in RPMI media with added 10% heat-inactivated fetal bovine serum (FBS) at 37°C in a 10% CO<sub>2</sub> atmosphere. Cells (approximately 1 x 10<sup>6</sup>  
10 cells per well for observation of markers; approximately 1 x 10<sup>5</sup> cells per well for detection of immunoglobulin in supernatant fluid) are cultured in the presence of 2.5 micrograms/ml of soluble, trimeric CD40L (described in the Armitage patents), for seven to eight days. Immunoglobulin concentrations (IgG, IgA) in the supernatant fluid are determined by specific ELISA; cells are monitored for the expression of CD54 by flow  
15 cytometry.

PBMC assays are conducted in RPMI 1640 complete medium. Approximately 2 x 10<sup>6</sup> cells per well are cultured in the presence of 2.5 micrograms/ml of soluble, trimeric CD40L (described in the Armitage patents), for 36 to 48 hours. Concentration of IL-12 and/or TNF-alpha in the supernatant fluid are determined by specific ELISA; cells are  
20 monitored for the expression of molecules indicative of activation by flow cytometry.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the  
25 foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A method for identifying compounds that alter one or more biological activities of CD40, comprising:
  - (a) mixing a test compound with a polypeptide selected from the group consisting of (i) a NEMO polypeptide comprising amino acids 287 through 419 SEQ ID NO:2, (ii) a fragment of a NEMO polypeptide according to (i) that is capable of binding a CYLD polypeptide according to SEQ ID NO:4 or fragment or variant thereof, and (iii) variants of the NEMO polypeptides of (i) and (ii); and
  - (b) determining whether the test compound alters the ability of NEMO to bind CYLD.
  
2. A method for identifying compounds that inhibit binding of NEMO and CYLD comprising:
  - (a) mixing a test compound with a polypeptide selected from the group consisting of (i) a NEMO polypeptide comprising amino acids 287 through 419 SEQ ID NO:2, (ii) a fragment of a NEMO polypeptide according to (i) that is capable of binding a CYLD polypeptide according to SEQ ID NO:4, and (iii) variants of the NEMO polypeptides of (i) and (ii), and a binding partner of said NEMO polypeptide selected from the group consisting of (iv) a CYLD polypeptide according to SEQ ID NO:4, (v) a fragment of a CYLD polypeptide according to SEQ ID NO:4 that is capable of binding a NEMO polypeptide of (i), (ii), or (iii), and (vi) a variant of the CYLD polypeptides of (iv) and (v) that is capable of binding a NEMO polypeptide of (i), (ii), or (iii); and
  - (b) determining whether the test compound inhibits the binding activity of said NEMO and CYLD polypeptides.
  
3. A method for producing information comprising the identity of a compound that alters one or more biological activities of CD40, the method comprising identifying one or more compounds that alter NEMO/CYLD binding activity.
  
4. The method of claim 3 wherein the compound decreases the binding of NEMO to CYLD.

5. The method of claim 3 wherein the compound increases the binding of NEMO to CYLD.

6. Information produced according to the method of claim 3, said information comprising the identity of a compound that alters the binding of NEMO to CYLD.

7. The information of claim 6 wherein the information is embodied in a storage medium selected from the group consisting of the brain of a living organism, paper, magnetic tape, optical tape, floppy disks, compact disks, computer system hard drives, and computer memory units.

8. A database comprising the information of claim 7.

9. The information of claim 7 wherein the information is embodied in a computer-readable medium.

10. The information of claim 7 wherein the information is embodied in a human-readable medium.

11. A computer system comprising a database containing records pertaining to a plurality of compounds, wherein the records comprise results of an assay according to claim 1, and a user interface allowing a user to access information regarding the plurality of compounds.

12. A computer system comprising a database containing records pertaining to a plurality of compounds, wherein the records comprise results of an assay according to claim 2, and a user interface allowing a user to access information regarding the plurality of compounds.

13. A computer system for storing and retrieving data on a plurality of compounds, the computer system comprising:

(a) input means for entering data for the compounds into a storage medium;

a processor for creating an individual record for each compound, the processor assigning specific identifying values for each compound;

(b) means for selecting one or more of the records based on results in an assay;

and

(c) means for transmitting information in the record or records to an output device to produce a report.

14. The system of claim 13 further comprising a video display unit.

15. The computer system of claim 13 where the report is in human-readable form.

16. The computer system of claim 14 where the report is in human-readable form.

17. A method of using the computer system of claim 13 to select one or more compounds for testing from a plurality of compounds having records stored in a database, the method comprising: displaying a list of said records or a field for entering information identifying one or more of said records; and selecting one or more of the records from the list or the record or records identified by entering information in the field.

18. A method of operating a computer system for analyzing compounds that modulate the interaction of NEMO and CYLD, the method comprising:

(a) entering data relating to a plurality of compounds into a storage medium; processing the data to create an individual record for each compound;

(b) testing compounds for the ability to modulate binding of NEMO to CYLD; and

(c) communicating results from the testing into the storage medium such that results for each compound are associated with the individual record for that compound:

19. The method of claim 18 wherein the storage medium comprises one or more computer memory units.

20. The method of claim 19 wherein the computer system further comprises a video display unit.

21. A database comprising records generated according to the method of claim 18.

22. A method of selecting compounds that alter one or more biological activities of CD40, comprising compiling a database according to claim 21, analyzing the testing results, and selecting one or more compounds.

23. A method for increasing one or more biological activities of CD40 comprising providing at least one compound selected according to the method of claim 22.

24. A method for decreasing one or more biological activities of CD40 comprising providing at least one compound selected according to the method of claim 22.




Figure 1



SEQUENCE LISTING

```

<110> IMMUNEX CORPORATION
      Derry, Jonathan M. J.
      Fanslow III, William
      Dougall, William C.

<120> SCREENING ASSAYS FOR AGONISTS OR ANTAGONISTS OF CD40 SIGNALING

<130> 3198-WO

<140> --to be assigned--
<141> 2002-05-07

<150> 09/851,673
<151> 2001-05-08

<160> 4

<170> PatentIn version 3.1

<210> 1
<211> 1994
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (149)..(1405)
<223>

<400> 1
ggcagcagca tggcccttgt gatccaggtg gggaaactaa ggcccagaga agtgaggacc      60
ccgcagacta tcaatcccag tctcttcccc tcaactccctg tgaagctctc cagcatcatc      120
gaggtcccat cagcccttgc cctgttgg atg aat agg cac ctc tgg aag agc          172
                Met Asn Arg His Leu Trp Lys Ser
                1                    5

caa ctg tgt gag atg gtg cag ccc agt ggt ggc ccg gca gca gat cag          220
Gln Leu Cys Glu Met Val Gln Pro Ser Gly Gly Pro Ala Ala Asp Gln
    10                    15                    20

gac gta ctg ggc gaa gag tct cct ctg ggg aag cca gcc atg ctg cac          268
Asp Val Leu Gly Glu Glu Ser Pro Leu Gly Lys Pro Ala Met Leu His
    25                    30                    35                    40

ctg cct tca gaa cag ggc gct cct gag acc ctc cag cgc tgc ctg gag          316
Leu Pro Ser Glu Gln Gly Ala Pro Glu Thr Leu Gln Arg Cys Leu Glu
                45                    50                    55

gag aat caa gag ctc cga gat gcc atc cgg cag agc aac cag att ctg          364
Glu Asn Gln Glu Leu Arg Asp Ala Ile Arg Gln Ser Asn Gln Ile Leu
    60                    65                    70
  
```

cgg gag cgc tgc gag gag ctt ctg cat ttc caa gcc agc cag agg gag Arg Glu Arg Cys Glu Glu Leu Leu His Phe Gln Ala Ser Gln Arg Glu	412
75 80 85	
gag aag gag ttc ctc atg tgc aag ttc cag gag gcc agg aaa ctg gtg Glu Lys Glu Phe Leu Met Cys Lys Phe Gln Glu Ala Arg Lys Leu Val	460
90 95 100	
gag aga ctc ggc ctg gag aag ctc gat ctg aag agg cag aag gag cag Glu Arg Leu Gly Leu Glu Lys Leu Asp Leu Lys Arg Gln Lys Glu Gln	508
105 110 115 120	
gct ctg cgg gag gtg gag cac ctg aag aga tgc cag cag cag atg gct Ala Leu Arg Glu Val Glu His Leu Lys Arg Cys Gln Gln Gln Met Ala	556
125 130 135	
gag gac aag gcc tct gtg aaa gcc cag gtg acg tcc ttg ctc ggg gag Glu Asp Lys Ala Ser Val Lys Ala Gln Val Thr Ser Leu Leu Gly Glu	604
140 145 150	
ctg cag gag agc cag agt cgc ttg gag gct gcc act aag gaa tgc cag Leu Gln Glu Ser Gln Ser Arg Leu Glu Ala Ala Thr Lys Glu Cys Gln	652
155 160 165	
gct ctg gag ggt cgg gcc cgg gcg gcc agc gag cag gcg cgg cag ctg Ala Leu Glu Gly Arg Ala Arg Ala Ala Ser Glu Gln Ala Arg Gln Leu	700
170 175 180	
gag agt gag cgc gag gcg ctg cag cag cag cac agc gtg cag gtg gac Glu Ser Glu Arg Glu Ala Leu Gln Gln Gln His Ser Val Gln Val Asp	748
185 190 195 200	
cag ctg cgc atg cag ggc cag agc gtg gag gcc gcg ctc cgc atg gag Gln Leu Arg Met Gln Gly Gln Ser Val Glu Ala Ala Leu Arg Met Glu	796
205 210 215	
cgc cag gcc gcc tcg gag gag aag agg aag ctg gcc cag ttg cag gtg Arg Gln Ala Ala Ser Glu Glu Lys Arg Lys Leu Ala Gln Leu Gln Val	844
220 225 230	
gcc tat cac cag ctc ttc caa gaa tac gac aac cac atc aag agc agc Ala Tyr His Gln Leu Phe Gln Glu Tyr Asp Asn His Ile Lys Ser Ser	892
235 240 245	
gtg gtg ggc agt gag cgg aag cga gga atg cag ctg gaa gat ctc aaa Val Val Gly Ser Glu Arg Lys Arg Gly Met Gln Leu Glu Asp Leu Lys	940
250 255 260	
cag cag ctc cag cag gcc gag gag gcc ctg gtg gcc aaa cag gag gtg Gln Gln Leu Gln Gln Ala Glu Glu Ala Leu Val Ala Lys Gln Glu Val	988
265 270 275 280	
atc gat aag ctg aag gag gag gcc gag cag cac aag att gtg atg gag Ile Asp Lys Leu Lys Glu Glu Ala Glu Gln His Lys Ile Val Met Glu	1036
285 290 295	

```

acc gtt ccg gtg ctg aag gcc cag gcg gat atc tac aag gcg gac ttc      1084
Thr Val Pro Val Leu Lys Ala Gln Ala Asp Ile Tyr Lys Ala Asp Phe
          300                      305                      310

cag gct gag agg cag gcc cgg gag aag ctg gcc gag aag aag gag ctc      1132
Gln Ala Glu Arg Gln Ala Arg Glu Lys Leu Ala Glu Lys Lys Glu Leu
          315                      320                      325

ctg cag gag cag ctg gag cag ctg cag agg gag tac agc aaa ctg aag      1180
Leu Gln Glu Gln Leu Glu Gln Leu Gln Arg Glu Tyr Ser Lys Leu Lys
          330                      335                      340

gcc agc tgt cag gag tgc gcc agg atc gag gac atg agg aag cgg cat      1228
Ala Ser Cys Gln Glu Ser Ala Arg Ile Glu Asp Met Arg Lys Arg His
          345                      350                      355                      360

gtc gag gtc tcc cag gcc ccc ttg ccc ccc gcc cct gcc tac ctc tcc      1276
Val Glu Val Ser Gln Ala Pro Leu Pro Pro Ala Pro Ala Tyr Leu Ser
          365                      370                      375

tct ccc ctg gcc ctg ccc agc cag agg agg agc ccc ccc gag gag cca      1324
Ser Pro Leu Ala Leu Pro Ser Gln Arg Arg Ser Pro Pro Glu Glu Pro
          380                      385                      390

cct gac ttc tgc tgt ccc aag tgc cag tat cag gcc cct gat atg gac      1372
Pro Asp Phe Cys Cys Pro Lys Cys Gln Tyr Gln Ala Pro Asp Met Asp
          395                      400                      405

acc ctg cag ata cat gtc atg gag tgc att gag tagggccggc cagtgcgaagg      1425
Thr Leu Gln Ile His Val Met Glu Cys Ile Glu
          410                      415

ccactgectg cccgaggacg tgcccgggac cgtgcagtct gcgctttcct ctcccgectg      1485

cctageccag gatgaagggc tgggtggcca caactgggat gccacctgga gccccacca      1545

ggagctggcc gcggcacctt acgcttcagc tgttgatccg ctggtcccet cttttgggggt      1605

agatgcgggc ccgatcaggc ctgactcgct gctctttttg ttcccttctg tctgctcgaa      1665

ccacttgect cgggctaate cctccctctt cctccaccgg gactgggga agtcaagaat      1725

ggggcctggg gctctcaggg agaactgctt ccctggcgag agctgggtgg cagctcttcc      1785

tcccaccgga caccgaccgg ccgcccgtg tgccctggga gtgctgcct cttaccatgc      1845

acacgggtgc tctccttttg ggctgcatgc tattccattt tgcagccaga ccgatgtgta      1905

tttaaccagt cactattgat ggacatttgg gttgtttccc atctttttgt taccataaat      1965

aatggcatag taaaaaaaaa aaaaaaaaaa      1994
    
```

<210> 2  
 <211> 419

<212> PRT

<213> Homo sapiens

<400> 2

Met Asn Arg His Leu Trp Lys Ser Gln Leu Cys Glu Met Val Gln Pro  
 1 5 10 15

Ser Gly Gly Pro Ala Ala Asp Gln Asp Val Leu Gly Glu Glu Ser Pro  
 20 25 30

Leu Gly Lys Pro Ala Met Leu His Leu Pro Ser Glu Gln Gly Ala Pro  
 35 40 45

Glu Thr Leu Gln Arg Cys Leu Glu Glu Asn Gln Glu Leu Arg Asp Ala  
 50 55 60

Ile Arg Gln Ser Asn Gln Ile Leu Arg Glu Arg Cys Glu Glu Leu Leu  
 65 70 75 80

His Phe Gln Ala Ser Gln Arg Glu Glu Lys Glu Phe Leu Met Cys Lys  
 85 90 95

Phe Gln Glu Ala Arg Lys Leu Val Glu Arg Leu Gly Leu Glu Lys Leu  
 100 105 110

Asp Leu Lys Arg Gln Lys Glu Gln Ala Leu Arg Glu Val Glu His Leu  
 115 120 125

Lys Arg Cys Gln Gln Gln Met Ala Glu Asp Lys Ala Ser Val Lys Ala  
 130 135 140

Gln Val Thr Ser Leu Leu Gly Glu Leu Gln Glu Ser Gln Ser Arg Leu  
 145 150 155 160

Glu Ala Ala Thr Lys Glu Cys Gln Ala Leu Glu Gly Arg Ala Arg Ala  
 165 170 175

Ala Ser Glu Gln Ala Arg Gln Leu Glu Ser Glu Arg Glu Ala Leu Gln  
 180 185 190

Gln Gln His Ser Val Gln Val Asp Gln Leu Arg Met Gln Gly Gln Ser  
 195 200 205

Val Glu Ala Ala Leu Arg Met Glu Arg Gln Ala Ala Ser Glu Glu Lys  
 210 215 220

Arg Lys Leu Ala Gln Leu Gln Val Ala Tyr His Gln Leu Phe Gln Glu  
 225 230 235 240

Tyr Asp Asn His Ile Lys Ser Ser Val Val Gly Ser Glu Arg Lys Arg  
 245 250 255

Gly Met Gln Leu Glu Asp Leu Lys Gln Gln Leu Gln Gln Ala Glu Glu  
 260 265 270

Ala Leu Val Ala Lys Gln Glu Val Ile Asp Lys Leu Lys Glu Glu Ala  
 275 280 285

Glu Gln His Lys Ile Val Met Glu Thr Val Pro Val Leu Lys Ala Gln  
 290 295 300

Ala Asp Ile Tyr Lys Ala Asp Phe Gln Ala Glu Arg Gln Ala Arg Glu  
 305 310 315 320

Lys Leu Ala Glu Lys Lys Glu Leu Leu Gln Glu Gln Leu Glu Gln Leu  
 325 330 335

Gln Arg Glu Tyr Ser Lys Leu Lys Ala Ser Cys Gln Glu Ser Ala Arg  
 340 345 350

Ile Glu Asp Met Arg Lys Arg His Val Glu Val Ser Gln Ala Pro Leu  
 355 360 365

Pro Pro Ala Pro Ala Tyr Leu Ser Ser Pro Leu Ala Leu Pro Ser Gln  
 370 375 380

Arg Arg Ser Pro Pro Glu Glu Pro Pro Asp Phe Cys Cys Pro Lys Cys  
 385 390 395 400

Gln Tyr Gln Ala Pro Asp Met Asp Thr Leu Gln Ile His Val Met Glu  
 405 410 415

Cys Ile Glu

<210> 3  
 <211> 5371  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (392)..(3262)  
 <223>

<400> 3

```

ggggggcgggc ccaggtagca ggtttggtg cgcgggggcc gcgcgtcgga gtttccccct      60
ttctaggggtg aggatggttc tacacagcca cccggagttc cttagttgaa aggtgcgccc      120
tgctgtgaca gaatgtggta attgtaatct ttaacatttt catgtaaac atatttctgt      180
atcatctttc cattgtcttc atggaaaatt gataaatatt tgtgccttcc aactctcgtc      240
ttggttgaat gacttcatct taatacaaca tggacaccac gttgctgaaa acatgctttg      300
ggactgccac tgaatttata ttttgcgggtt ttatgacaaa gttattagta gtttcccttt      360
tttgaattag tattttgaag ttaatatac a atg agt tca ggc tta tgg agc      412
                               Met Ser Ser Gly Leu Trp Ser
                               1                               5

caa gaa aaa gtc act tca ccc tac tgg gaa gag cgg att ttt tac ttg      460
Gln Glu Lys Val Thr Ser Pro Tyr Trp Glu Glu Arg Ile Phe Tyr Leu
                               10                               15                               20

ctt ctt caa gaa tgc agc gtt aca gac aaa caa aca caa aag ctc ctt      508
Leu Leu Gln Glu Cys Ser Val Thr Asp Lys Gln Thr Gln Lys Leu Leu
                               25                               30                               35

aaa gta ccg aag gga agt ata gga cag tat att caa gat cgt tct gtg      556
Lys Val Pro Lys Gly Ser Ile Gly Gln Tyr Ile Gln Asp Arg Ser Val
40                               45                               50                               55

ggg cat tca agg att cct tct gca aaa ggc aag aaa aat cag att gga      604
Gly His Ser Arg Ile Pro Ser Ala Lys Gly Lys Lys Asn Gln Ile Gly
                               60                               65                               70

tta aaa att cta gag caa cct cat gca gtt ctc ttt gtt gat gaa aag      652
Leu Lys Ile Leu Glu Gln Pro His Ala Val Leu Phe Val Asp Glu Lys
                               75                               80                               85

gat gtt gta gag ata aat gaa aag ttc aca gag tta ctt ttg gca att      700
Asp Val Val Glu Ile Asn Glu Lys Phe Thr Glu Leu Leu Leu Ala Ile
                               90                               95                               100

acc aat tgt gag gag agg ttc agc ctg ttt aaa aac aga aac aga cta      748
Thr Asn Cys Glu Glu Arg Phe Ser Leu Phe Lys Asn Arg Asn Arg Leu
105                               110                               115

```

agt	aaa	ggc	ctc	caa	ata	gac	gtg	ggc	tgt	cct	gtg	aaa	gta	cag	ctg	796
Ser	Lys	Gly	Leu	Gln	Ile	Asp	Val	Gly	Cys	Pro	Val	Lys	Val	Gln	Leu	
120					125					130					135	
aga	tct	ggg	gaa	gaa	aaa	ttt	cct	gga	ggt	gta	cgc	ttc	aga	gga	ccc	844
Arg	Ser	Gly	Glu	Glu	Lys	Phe	Pro	Gly	Val	Val	Arg	Phe	Arg	Gly	Pro	
			140						145						150	
ctg	tta	gca	gag	agg	aca	gtc	tcc	gga	ata	ttc	ttt	gga	ggt	gaa	ttg	892
Leu	Leu	Ala	Glu	Arg	Thr	Val	Ser	Gly	Ile	Phe	Phe	Gly	Val	Glu	Leu	
			155					160					165			
ctg	gaa	gaa	ggg	cgt	ggg	caa	ggg	ttc	act	gac	ggg	gtg	tac	caa	ggg	940
Leu	Glu	Glu	Gly	Arg	Gly	Gln	Gly	Phe	Thr	Asp	Gly	Val	Tyr	Gln	Gly	
	170						175					180				
aaa	cag	ctt	ttt	cag	tgt	gat	gaa	gat	tgt	ggc	gtg	ttt	ggt	gca	ttg	988
Lys	Gln	Leu	Phe	Gln	Cys	Asp	Glu	Asp	Cys	Gly	Val	Phe	Val	Ala	Leu	
	185					190					195					
gac	aag	cta	gaa	ctc	ata	gaa	gat	gat	gac	act	gca	ttg	gaa	agt	gat	1036
Asp	Lys	Leu	Glu	Leu	Ile	Glu	Asp	Asp	Asp	Thr	Ala	Leu	Glu	Ser	Asp	
200					205					210					215	
tac	gca	ggg	cct	ggg	gac	aca	atg	cag	gtc	gaa	ctt	cct	cct	ttg	gaa	1084
Tyr	Ala	Gly	Pro	Gly	Asp	Thr	Met	Gln	Val	Glu	Leu	Pro	Pro	Leu	Glu	
				220					225						230	
ata	aac	tcc	aga	ggt	tct	ttg	aag	ggt	gga	gaa	aca	ata	gaa	tct	gga	1132
Ile	Asn	Ser	Arg	Val	Ser	Leu	Lys	Val	Gly	Glu	Thr	Ile	Glu	Ser	Gly	
			235					240							245	
aca	ggt	ata	ttc	tgt	gat	ggt	ttg	cca	gga	aaa	gaa	agc	tta	gga	tat	1180
Thr	Val	Ile	Phe	Cys	Asp	Val	Leu	Pro	Gly	Lys	Glu	Ser	Leu	Gly	Tyr	
		250					255					260				
ttt	ggt	ggg	gtg	gac	atg	gat	aac	cct	att	ggc	aac	tgg	gat	gga	aga	1228
Phe	Val	Gly	Val	Asp	Met	Asp	Asn	Pro	Ile	Gly	Asn	Trp	Asp	Gly	Arg	
	265					270					275					
ttt	gat	gga	gtg	cag	ctt	tgt	agt	ttt	gcg	tgt	ggt	gaa	agt	aca	att	1276
Phe	Asp	Gly	Val	Gln	Leu	Cys	Ser	Phe	Ala	Cys	Val	Glu	Ser	Thr	Ile	
280					285					290					295	
cta	ttg	cac	atc	aat	gat	atc	atc	cca	gct	tta	tca	gag	agt	gtg	acg	1324
Leu	Leu	His	Ile	Asn	Asp	Ile	Ile	Pro	Ala	Leu	Ser	Glu	Ser	Val	Thr	
				300					305						310	
cag	gaa	agg	agg	cct	ccc	aaa	ctt	gcc	ttt	atg	tca	aga	ggg	ggt	ggg	1372
Gln	Glu	Arg	Arg	Pro	Pro	Lys	Leu	Ala	Phe	Met	Ser	Arg	Gly	Val	Gly	
			315					320							325	
gac	aaa	ggg	tca	tcc	agt	cat	aat	aaa	cca	aag	gct	aca	gga	tct	acc	1420
Asp	Lys	Gly	Ser	Ser	Ser	His	Asn	Lys	Pro	Lys	Ala	Thr	Gly	Ser	Thr	
		330					335					340				

tca gac cct gga aat aga aac aga tct gaa tta ttt tat acc tta aat	1468
Ser Asp Pro Gly Asn Arg Asn Arg Ser Glu Leu Phe Tyr Thr Leu Asn	
345 350 355	
ggg tct tct gtt gac tca caa cca caa tcc aaa tca aaa aat aca tgg	1516
Gly Ser Ser Val Asp Ser Gln Pro Gln Ser Lys Ser Lys Asn Thr Trp	
360 365 370 375	
tac att gat gaa gtt gca gaa gac cct gca aaa tct ctt aca gag ata	1564
Tyr Ile Asp Glu Val Ala Glu Asp Pro Ala Lys Ser Leu Thr Glu Ile	
380 385 390	
tct aca gac ttt gac cgt tct tca cca cca ctc cag cct cct cct gtg	1612
Ser Thr Asp Phe Asp Arg Ser Ser Pro Pro Leu Gln Pro Pro Pro Val	
395 400 405	
aac tca ctg acc acc gag aac aga ttc cac tct tta cca ttc agt ctc	1660
Asn Ser Leu Thr Thr Glu Asn Arg Phe His Ser Leu Pro Phe Ser Leu	
410 415 420	
acc aag atg ccc aat acc aat gga agt att ggc cac agt cca ctt tct	1708
Thr Lys Met Pro Asn Thr Asn Gly Ser Ile Gly His Ser Pro Leu Ser	
425 430 435	
ctg tca gcc cag tct gta atg gaa gag cta aac act gca ccc gtc caa	1756
Leu Ser Ala Gln Ser Val Met Glu Glu Leu Asn Thr Ala Pro Val Gln	
440 445 450 455	
gag agt cca ccc ttg gcc atg cct cct ggg aac tca cat ggt cta gaa	1804
Glu Ser Pro Pro Leu Ala Met Pro Pro Gly Asn Ser His Gly Leu Glu	
460 465 470	
gtg ggc tca ttg gct gaa gtt aag gag aac cct cct ttc tat ggg gta	1852
Val Gly Ser Leu Ala Glu Val Lys Glu Asn Pro Pro Phe Tyr Gly Val	
475 480 485	
atc cgt tgg atc ggt cag cca cca gga ctg aat gaa gtg ctc gct gga	1900
Ile Arg Trp Ile Gly Gln Pro Pro Gly Leu Asn Glu Val Leu Ala Gly	
490 495 500	
ctg gaa ctg gaa gat gag tgt gca ggc tgt acg gat gga acc ttc aga	1948
Leu Glu Leu Glu Asp Glu Cys Ala Gly Cys Thr Asp Gly Thr Phe Arg	
505 510 515	
ggc act cgg tat ttc acc tgt gcc ctg aag aag gcg ctg ttt gtg aaa	1996
Gly Thr Arg Tyr Phe Thr Cys Ala Leu Lys Lys Ala Leu Phe Val Lys	
520 525 530 535	
ctg aag agc tgc agg cct gac tct agg ttt gca tca ttg cag ccg gtt	2044
Leu Lys Ser Cys Arg Pro Asp Ser Arg Phe Ala Ser Leu Gln Pro Val	
540 545 550	
tcc aat cag att gag cgc tgt aac tct tta gca ttt gga ggc tac tta	2092
Ser Asn Gln Ile Glu Arg Cys Asn Ser Leu Ala Phe Gly Gly Tyr Leu	
555 560 565	



agt	gaa	gta	gta	gaa	gaa	aat	act	cca	cca	aaa	atg	gaa	aaa	gaa	ggc	2140
Ser	Glu	Val	Val	Glu	Glu	Asn	Thr	Pro	Pro	Lys	Met	Glu	Lys	Glu	Gly	
		570					575					580				
ttg	gag	ata	atg	att	ggg	aag	aag	aaa	ggc	atc	cag	ggt	cat	tac	aat	2188
Leu	Glu	Ile	Met	Ile	Gly	Lys	Lys	Lys	Gly	Ile	Gln	Gly	His	Tyr	Asn	
	585					590				595						
tct	tgt	tac	tta	gac	tca	acc	tta	ttc	tgc	tta	ttt	gct	ttt	agt	tct	2236
Ser	Cys	Tyr	Leu	Asp	Ser	Thr	Leu	Phe	Cys	Leu	Phe	Ala	Phe	Ser	Ser	
600				605					610					615		
gtt	ctg	gac	act	gtg	tta	ctt	aga	ccc	aaa	gaa	aag	aac	gat	gta	gaa	2284
Val	Leu	Asp	Thr	Val	Leu	Leu	Arg	Pro	Lys	Glu	Lys	Asn	Asp	Val	Glu	
			620					625					630			
tat	tat	agt	gaa	acc	caa	gag	cta	ctg	agg	aca	gaa	att	gtt	aat	cct	2332
Tyr	Tyr	Ser	Glu	Thr	Gln	Glu	Leu	Leu	Arg	Thr	Glu	Ile	Val	Asn	Pro	
		635				640						645				
ctg	aga	ata	tat	gga	tat	gtg	tgt	gcc	aca	aaa	att	atg	aaa	ctg	agg	2380
Leu	Arg	Ile	Tyr	Gly	Tyr	Val	Cys	Ala	Thr	Lys	Ile	Met	Lys	Leu	Arg	
	650			655				660								
aaa	ata	ctt	gaa	aag	gtg	gag	gct	gca	tca	gga	ttt	acc	tct	gaa	gaa	2428
Lys	Ile	Leu	Glu	Lys	Val	Glu	Ala	Ala	Ser	Gly	Phe	Thr	Ser	Glu	Glu	
	665			670				675								
aaa	gat	cct	gag	gaa	ttc	ttg	aat	att	ctg	ttt	cat	cat	att	tta	agg	2476
Lys	Asp	Pro	Glu	Glu	Phe	Leu	Asn	Ile	Leu	Phe	His	His	Ile	Leu	Arg	
680			685					690						695		
gta	gaa	cct	ttg	cta	aaa	ata	aga	tca	gca	ggt	caa	aag	gta	caa	gat	2524
Val	Glu	Pro	Leu	Leu	Lys	Ile	Arg	Ser	Ala	Gly	Gln	Lys	Val	Gln	Asp	
			700					705					710			
tgt	tac	ttc	tat	caa	att	ttt	atg	gaa	aaa	aat	gag	aaa	gtt	ggc	gtt	2572
Cys	Tyr	Phe	Tyr	Gln	Ile	Phe	Met	Glu	Lys	Asn	Glu	Lys	Val	Gly	Val	
		715				720							725			
ccc	aca	att	cag	cag	ttg	tta	gaa	tgg	tct	ttt	atc	aac	agt	aac	ctg	2620
Pro	Thr	Ile	Gln	Gln	Leu	Leu	Glu	Trp	Ser	Phe	Ile	Asn	Ser	Asn	Leu	
		730				735						740				
aaa	ttt	gca	gag	gca	cca	tca	tgt	ctg	att	att	cag	atg	cct	cga	ttt	2668
Lys	Phe	Ala	Glu	Ala	Pro	Ser	Cys	Leu	Ile	Ile	Gln	Met	Pro	Arg	Phe	
	745					750					755					
gga	aaa	gac	ttt	aaa	cta	ttt	aaa	aaa	att	ttt	cct	tct	ctg	gaa	tta	2716
Gly	Lys	Asp	Phe	Lys	Leu	Phe	Lys	Lys	Ile	Phe	Pro	Ser	Leu	Glu	Leu	
760				765				770						775		
aat	ata	aca	gat	tta	ctt	gaa	gac	act	ccc	aga	cag	tgc	cgg	ata	tgt	2764
Asn	Ile	Thr	Asp	Leu	Leu	Glu	Asp	Thr	Pro	Arg	Gln	Cys	Arg	Ile	Cys	
			780					785						790		

gga ggg ctt gca atg tat gag tgt aga gaa tgc tac gac gat ccg gac 2812  
 Gly Gly Leu Ala Met Tyr Glu Cys Arg Glu Cys Tyr Asp Asp Pro Asp  
 795 800 805

atc tca gct gga aaa atc aag cag ttt tgt aaa acc tgc aac act caa 2860  
 Ile Ser Ala Gly Lys Ile Lys Gln Phe Cys Lys Thr Cys Asn Thr Gln  
 810 815 820

gtc cac ctt cat ccg aag agg ctg aat cat aaa tat aac cca gtg tca 2908  
 Val His Leu His Pro Lys Arg Leu Asn His Lys Tyr Asn Pro Val Ser  
 825 830 835

ctt ccc aaa gac tta ccc gac tgg gac tgg aga cac ggc tgc atc cct 2956  
 Leu Pro Lys Asp Leu Pro Asp Trp Asp Trp Arg His Gly Cys Ile Pro  
 840 845 850 855

tgc cag aat atg gag tta ttt gct gtt ctc tgc ata gaa aca agc cac 3004  
 Cys Gln Asn Met Glu Leu Phe Ala Val Leu Cys Ile Glu Thr Ser His  
 860 865 870

tat gtt gct ttt gtg aag tat ggg aag gac gat tct gcc tgg ctc ttc 3052  
 Tyr Val Ala Phe Val Lys Tyr Gly Lys Asp Asp Ser Ala Trp Leu Phe  
 875 880 885

ttt gac agc atg gcc gat cgg gat ggt ggt cag aat ggc ttc aac att 3100  
 Phe Asp Ser Met Ala Asp Arg Asp Gly Gly Gln Asn Gly Phe Asn Ile  
 890 895 900

cct caa gtc acc cca tgc cca gaa gta gga gag tac ttg aag atg tct 3148  
 Pro Gln Val Thr Pro Cys Pro Glu Val Gly Glu Tyr Leu Lys Met Ser  
 905 910 915

ctg gaa gac ctg cat tcc ttg gac tcc agg aga atc caa ggc tgt gca 3196  
 Leu Glu Asp Leu His Ser Leu Asp Ser Arg Arg Ile Gln Gly Cys Ala  
 920 925 930 935

cga aga ctg ctt tgt gat gca tat atg tgc atg tac cag agt cca aca 3244  
 Arg Arg Leu Leu Cys Asp Ala Tyr Met Cys Met Tyr Gln Ser Pro Thr  
 940 945 950

atg agt ttg tac aaa taa ctggggtcat cgggaaaggc aaagaaactg 3292  
 Met Ser Leu Tyr Lys  
 955

aaggcagagt cctaacgttg catcttattc gagctggcag ttctgttcac gtccattgcc 3352

ggcaatggat gtctttgtgg tgatgatcct tcagaaaagg atgcctctgt ttaaaaacaa 3412

attgcttttg tgtccctgaa gtatttaata agaagcattt tgcactctag aaagtatggt 3472

tgtgttggtt ttttaagaag tctaaatgaa gttattaata cctgaagctt taagttaagt 3532

gcattgatca tatgatattt ttggaagcat acaattttaa ttgtggaagt ttaaagcctc 3592

ttttagtcca ttgagaatgt aaataaatgt gtcttcttta tggaccaagg atatgaaac 3652

atTTTTcttt	tgtagctaac	ggttgccttg	aggaagaaat	aatttggttt	tattaagagt	3712
ctactctcaa	tccagttatt	agagatgtac	tgagtttgat	ttgttaatcc	tttctatata	3772
ctgctgatct	tgcattgtcta	caatctgctc	agtttttctg	tgtttctgca	atagtgggtca	3832
gaaaaaact	taaattccct	taatgggtgt	gttttctatt	tgttctgggt	ttgagataaa	3892
tgagtgatc	tgtcccaaaa	tgtccatttt	tgaagtgatt	ttcctggagg	attaggggat	3952
ttagcagttg	aagctcttca	ttcatagtag	ttactgtcag	ctaacagggt	ttttaaggct	4012
tttaactatt	aatattttat	ggaatggggc	aaagtaaatt	gatgaaagaa	ttggagtgat	4072
aatagtcctt	tacaaacata	cagtccataa	gaaaatgaat	ttggcatata	gaattattac	4132
aatttcctgg	gagagatgga	tatttaaaoc	tctattattt	tagacaagac	tgtctagaac	4192
ttaagtttga	tctgtcagcc	agtactcca	ttaaattcag	tgtagtttca	cttgatagaa	4252
tcagatatgt	tatcgaaatg	ttagcagcag	cttcatcctc	cttctgatta	aagtaagtag	4312
aatgggatg	ttttgtttaa	taacagccat	agtgtgtgtt	tagaccacag	cggatgttgt	4372
agaccaggac	catagatgat	acatgtcagt	gctgtggaat	gtgcattctc	tgagtgttgt	4432
tttgtggtat	cattgtcttt	cctgaatgac	tttctaactg	tgagaaaagg	cagaaaagtc	4492
atcatatgta	tatgtcatat	gactttataa	aatatttaat	gtgacaaaaa	gtggaaagaa	4552
tctttacaaa	ccctgcaatt	acttttttaa	aggcactttt	actctttggt	tttatcattc	4612
cattttgcta	atatttacta	gctttataaa	ttacagtaag	gtacaaaaac	tcattcttgta	4672
atattttcat	ttttgaagtg	aaaaagtaca	tatattttgc	acaaggtttt	atactgctaa	4732
gtgcttggtt	ggggtggtga	gatgatgatt	agatcagggg	tgaggctgag	agactctggg	4792
tttagggcta	gcctgcctc	catctccctt	gggtaaaatg	aagggtgtgg	ggtaaaagat	4852
gcataaggcc	ttttctagct	ctgacagcct	agaagtcaa	tcaccctgta	ataaatatgt	4912
gttgaatgaa	gaaatgggtg	aatgagcttg	tcaatgtgat	tttaaaaaat	tgactacctg	4972
gaggaatgat	taggaatcta	aatgaagcca	gccctcggtg	tctgcagggt	tctcatccat	5032
ggattcaacc	aactgcaaat	ggaaaatagc	atTTTTTTta	aaaaaaggat	ggttacatcc	5092
gtattgaaca	tgtacagact	tttttcttgt	cattattctc	tgaacaatac	agaactctt	5152
tatgtagcat	ttacatttat	taggtattat	aagtaatcta	gagattattt	aattaaaata	5212
tacaggagga	tgtgtgttta	tatgccagaa	attctgtacc	atTTTgtatc	agggatttga	5272
gcatcttcag	atggttggtat	ctgcagggat	cctggaacca	aaccctgca	gataactaagg	5332

gctgacgatc taggtaagac tggatttaac agttggaaa

5371

<210> 4  
 <211> 956  
 <212> PRT  
 <213> Homo sapiens

<400> 4

Met Ser Ser Gly Leu Trp Ser Gln Glu Lys Val Thr Ser Pro Tyr Trp  
 1 5 10 15

Glu Glu Arg Ile Phe Tyr Leu Leu Leu Gln Glu Cys Ser Val Thr Asp  
 20 25 30

Lys Gln Thr Gln Lys Leu Leu Lys Val Pro Lys Gly Ser Ile Gly Gln  
 35 40 45

Tyr Ile Gln Asp Arg Ser Val Gly His Ser Arg Ile Pro Ser Ala Lys  
 50 55 60

Gly Lys Lys Asn Gln Ile Gly Leu Lys Ile Leu Glu Gln Pro His Ala  
 65 70 75 80

Val Leu Phe Val Asp Glu Lys Asp Val Val Glu Ile Asn Glu Lys Phe  
 85 90 95

Thr Glu Leu Leu Leu Ala Ile Thr Asn Cys Glu Glu Arg Phe Ser Leu  
 100 105 110

Phe Lys Asn Arg Asn Arg Leu Ser Lys Gly Leu Gln Ile Asp Val Gly  
 115 120 125

Cys Pro Val Lys Val Gln Leu Arg Ser Gly Glu Glu Lys Phe Pro Gly  
 130 135 140

Val Val Arg Phe Arg Gly Pro Leu Leu Ala Glu Arg Thr Val Ser Gly  
 145 150 155 160

Ile Phe Phe Gly Val Glu Leu Leu Glu Glu Gly Arg Gly Gln Gly Phe  
 165 170 175

Thr Asp Gly Val Tyr Gln Gly Lys Gln Leu Phe Gln Cys Asp Glu Asp  
 180 185 190

Cys Gly Val Phe Val Ala Leu Asp Lys Leu Glu Leu Ile Glu Asp Asp  
 195 200 205

Asp Thr Ala Leu Glu Ser Asp Tyr Ala Gly Pro Gly Asp Thr Met Gln  
 210 215 220

Val Glu Leu Pro Pro Leu Glu Ile Asn Ser Arg Val Ser Leu Lys Val  
 225 230 235 240

Gly Glu Thr Ile Glu Ser Gly Thr Val Ile Phe Cys Asp Val Leu Pro  
 245 250 255

Gly Lys Glu Ser Leu Gly Tyr Phe Val Gly Val Asp Met Asp Asn Pro  
 260 265 270

Ile Gly Asn Trp Asp Gly Arg Phe Asp Gly Val Gln Leu Cys Ser Phe  
 275 280 285

Ala Cys Val Glu Ser Thr Ile Leu Leu His Ile Asn Asp Ile Ile Pro  
 290 295 300

Ala Leu Ser Glu Ser Val Thr Gln Glu Arg Arg Pro Pro Lys Leu Ala  
 305 310 315 320

Phe Met Ser Arg Gly Val Gly Asp Lys Gly Ser Ser Ser His Asn Lys  
 325 330 335

Pro Lys Ala Thr Gly Ser Thr Ser Asp Pro Gly Asn Arg Asn Arg Ser  
 340 345 350

Glu Leu Phe Tyr Thr Leu Asn Gly Ser Ser Val Asp Ser Gln Pro Gln  
 355 360 365

Ser Lys Ser Lys Asn Thr Trp Tyr Ile Asp Glu Val Ala Glu Asp Pro  
 370 375 380

Ala Lys Ser Leu Thr Glu Ile Ser Thr Asp Phe Asp Arg Ser Ser Pro  
 385 390 395 400

Pro Leu Gln Pro Pro Pro Val Asn Ser Leu Thr Thr Glu Asn Arg Phe  
 405 410 415

His Ser Leu Pro Phe Ser Leu Thr Lys Met Pro Asn Thr Asn Gly Ser  
 420 425 430

Ile Gly His Ser Pro Leu Ser Leu Ser Ala Gln Ser Val Met Glu Glu  
 435 440 445

Leu Asn Thr Ala Pro Val Gln Glu Ser Pro Pro Leu Ala Met Pro Pro  
 450 455 460

Gly Asn Ser His Gly Leu Glu Val Gly Ser Leu Ala Glu Val Lys Glu  
 465 470 475 480

Asn Pro Pro Phe Tyr Gly Val Ile Arg Trp Ile Gly Gln Pro Pro Gly  
 485 490 495

Leu Asn Glu Val Leu Ala Gly Leu Glu Leu Glu Asp Glu Cys Ala Gly  
 500 505 510

Cys Thr Asp Gly Thr Phe Arg Gly Thr Arg Tyr Phe Thr Cys Ala Leu  
 515 520 525

Lys Lys Ala Leu Phe Val Lys Leu Lys Ser Cys Arg Pro Asp Ser Arg  
 530 535 540

Phe Ala Ser Leu Gln Pro Val Ser Asn Gln Ile Glu Arg Cys Asn Ser  
 545 550 555 560 565

Leu Ala Phe Gly Gly Tyr Leu Ser Glu Val Val Glu Glu Asn Thr Pro  
 565 570 575

Pro Lys Met Glu Lys Glu Gly Leu Glu Ile Met Ile Gly Lys Lys Lys  
 580 585 590

Gly Ile Gln Gly His Tyr Asn Ser Cys Tyr Leu Asp Ser Thr Leu Phe  
 595 600 605

Cys Leu Phe Ala Phe Ser Ser Val Leu Asp Thr Val Leu Leu Arg Pro  
 610 615 620

Lys Glu Lys Asn Asp Val Glu Tyr Tyr Ser Glu Thr Gln Glu Leu Leu  
625 630 635 640

Arg Thr Glu Ile Val Asn Pro Leu Arg Ile Tyr Gly Tyr Val Cys Ala  
645 650 655

Thr Lys Ile Met Lys Leu Arg Lys Ile Leu Glu Lys Val Glu Ala Ala  
660 665 670

Ser Gly Phe Thr Ser Glu Glu Lys Asp Pro Glu Glu Phe Leu Asn Ile  
675 680 685

Leu Phe His His Ile Leu Arg Val Glu Pro Leu Leu Lys Ile Arg Ser  
690 695 700

Ala Gly Gln Lys Val Gln Asp Cys Tyr Phe Tyr Gln Ile Phe Met Glu  
705 710 715 720

Lys Asn Glu Lys Val Gly Val Pro Thr Ile Gln Gln Leu Leu Glu Trp  
725 730 735

Ser Phe Ile Asn Ser Asn Leu Lys Phe Ala Glu Ala Pro Ser Cys Leu  
740 745 750

Ile Ile Gln Met Pro Arg Phe Gly Lys Asp Phe Lys Leu Phe Lys Lys  
755 760 765

Ile Phe Pro Ser Leu Glu Leu Asn Ile Thr Asp Leu Leu Glu Asp Thr  
770 775 780

Pro Arg Gln Cys Arg Ile Cys Gly Gly Leu Ala Met Tyr Glu Cys Arg  
785 790 795 800

Glu Cys Tyr Asp Asp Pro Asp Ile Ser Ala Gly Lys Ile Lys Gln Phe  
805 810 815

Cys Lys Thr Cys Asn Thr Gln Val His Leu His Pro Lys Arg Leu Asn  
820 825 830

His Lys Tyr Asn Pro Val Ser Leu Pro Lys Asp Leu Pro Asp Trp Asp  
835 840 845

Trp Arg His Gly Cys Ile Pro Cys Gln Asn Met Glu Leu Phe Ala Val  
850 855 860

Leu Cys Ile Glu Thr Ser His Tyr Val Ala Phe Val Lys Tyr Gly Lys  
865 870 875 880

Asp Asp Ser Ala Trp Leu Phe Phe Asp Ser Met Ala Asp Arg Asp Gly  
885 890 895

Gly Gln Asn Gly Phe Asn Ile Pro Gln Val Thr Pro Cys Pro Glu Val  
900 905 910

Gly Glu Tyr Leu Lys Met Ser Leu Glu Asp Leu His Ser Leu Asp Ser  
915 920 925

Arg Arg Ile Gln Gly Cys Ala Arg Arg Leu Leu Cys Asp Ala Tyr Met  
930 935 940

Cys Met Tyr Gln Ser Pro Thr Met Ser Leu Tyr Lys  
945 950 955